

※ : Revised in March 2023

- **This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.**
- **Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.**
- **Multilingual product information is a tentative translation by the provider (or Our Company), and may be modified or altered without notice.**
- **The provider (or Our Company) assumes no responsibility for any problems that occur due to the contents of the multilingual product information.**

SEIROGAN
Quick C

You should read this written explanation before use.
Please keep it so that you can read it when you need to.



大幸薬品

正露丸クイック C

2nd Class OTC drug

Gastrointestinal
medicine

Seirogan Quick C

Seirogan Quick C is a liquid capsule that rapidly dissolves JP wood creosote, which is a crude drug contained in Seirogan, a household medicine used for over a century. Japanese Pharmacopoeia (JP) wood creosote is a clear liquid extracted from trees such as beech and pine which regulates the water balance in the intestines without stopping bowel peristalsis, and returns the intestinal environment to its normal condition. Against the sudden diarrhea and soft stool that can occur when commuting, going to school or going out, we recommend taking prompt action with **Seirogan Quick C**.



[Precautions]

Consultation



1. The following persons should contact a physician, pharmacist, or registered salesperson for a consultation before administration.
 - (1) Patients undergoing medical treatment from a physician
 - (2) Those who have diarrhea with fever, bloody stool or persistent mucous stool
 - (3) The elderly (age-related decline of kidney and liver function may amplify the effect of drugs)
2. If the following symptoms are observed after using this drug, these may be adverse reactions, so immediately discontinue the use of this drug, and show this document (package insert) to your physician, dentist, pharmacist, or registered salesperson for a consultation.

Related area	Symptom
Gastrointestinal system	Loss of appetite, feeling of discomfort in the stomach

(Overdose of JP wood creosote, the main active ingredient, may cause gastric disorders and loss of appetite.)

3. When symptoms do not improve even after taking this medicine for 5 to 6 days, stop taking this medicine and consult a physician, dentist, pharmacist or registered salesperson, being sure to take this package insert (instruction leaflet) with you.

(Continued overleaf)

[Indications]

Diarrhea, diarrhea due to indigestion, food poisoning, water poisoning, diarrhea and vomiting, loose bowels and soft stool

[Dosage and directions]

Take the following dosages no more than 3 times a day with cold or warm water with an interval of at least 4 hours between doses.

Age	One dose	Daily dosage
11 years and over	2 capsules	No more than 3 times
5-10 years of age	1 capsule	
Under 5 years of age	Do not use	

[Precautions of Dosage and Administration]

- (1) Follow the dosage and directions.
- (2) When allowing children 5 years or over to take the medicine, a guardian must be present, and caution must be exercised to prevent them from taking an incorrect dose or choking on it.
- (3) To take a capsule from the package: As shown in the illustration on the right, press the convex part of the blister pack sheet containing the capsule firmly with a fingertip, break the aluminum foil on the back side, and take out the capsule. (Accidental ingestion of a blister pack sheet containing the capsule can lead to injury, such as damage to the esophageal membrane).



[Ingredients and contents]

Every 2 capsules (single dosage of 11 years and over) include the following ingredients:

Active Ingredient	Contents	Effect
JP wood creosote	90 mg	Normalization of excessive colonic motility, regulation of intestinal water content, bacteriostasis inside the intestines

Inactive ingredients: Macrogol 400, polyoxyethylene 80, succinylated gelatin, D-sorbitol, glycerin, caramel and medium chain fatty acid triglyceride.

[Precautions for storage and handling]

1. Keep out of reach of children.
2. Do not transfer the medicine to a different container. (It may lead to misuse or alter the quality of the drug.)
3. Once the package is opened, fold the aluminum pouch at the folding lines and always store the product in the aluminum pouch together with desiccant.
4. Store in a cool place away from direct sunlight.
5. Do not use after the use-by date indicated on the package.

For more information, please visit our website (<http://www.seirogan.co.jp/>).

Marketing authorization holder

TAIKO PHARMACEUTICAL CO., LTD.

3-34-14, Uchihonmachi, Suita Osaka, Japan 564-0032